

CLAIMS

We claim:

1. An array composition comprising:

a) a substrate with a surface comprising discrete sites; and

b) a population of microspheres comprising at least a first and a second subpopulation,

wherein the microspheres of each subpopulation each comprise a plurality of target analytes;

wherein said microspheres are distributed on said surface.

2. The array composition according to claim 1 wherein said microspheres of each subpopulation further comprise an optical signature.

3. The array composition according to claim 1 wherein said microspheres of each subpopulation further comprise an identifier binding ligand.

4. The array composition according to claim 3 wherein said identifier binding ligand is a nucleic acid.

5. The array composition according to claim 1 wherein said target analytes are nucleic acids.

6. The array composition according to claim 5 wherein said nucleic acids comprise genomic DNA.

7. The array composition according to claim 1 wherein said target analytes are proteins.

8. The array composition according to claim 1 wherein said substrate is a fiber optic substrate.

9. The array composition according to claim 1 wherein said substrate is plastic.

10. The array composition according to claim 1 wherein said discrete sites are wells.

11. The array composition according to claim 1, wherein said microspheres are randomly distributed on said surface.

12. The array composition according to claim 1, wherein the microspheres of said first and second subpopulation each comprise a plurality of target analytes from a first and second target source, respectively.

13. The array composition according to claim 13, wherein said first and second target source are first and second patients, respectively.

14. A method comprising:

a) providing an array composition comprising:

- i) a substrate with a surface comprising discrete sites; and
- ii) a population of microspheres comprising at least a first and a second subpopulation, wherein the microspheres of each subpopulation each comprise a plurality of target analytes;

wherein said microspheres are distributed on said surface;

b) contacting said array composition with a first set of readout probes;

c) detecting the presence of a first target analyte.

15. The method according to claim 14 further comprising:

d) contacting said array composition with a second set of readout probes;

e) detecting the presence of a second target analyte.

16. The method according to claim 14, wherein said microspheres are randomly distributed on said surface.

17. The method according to claim 14, wherein said first set of readout probes comprises at least first and second readout probes, wherein said first and second readout probes comprise first and second labels, respectively.

18. The method according to claim 17, further comprising detecting said first label as an indication of the presence of said first target analyte.

19. The method according to claim 14, wherein the microspheres of said first and second subpopulation each comprise a plurality of target analytes from a first and second target source, respectively.

20. The array composition according to claim 19, wherein said first and second target source are first and second patients, respectively.

21. A method of genotyping comprising:

a) providing an array composition comprising:

5  
Sub B1

10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100

- 5
- Sub B1
- i) a substrate with a surface comprising discrete sites; and
  - ii) a population of microspheres comprising at least a first and a second subpopulation, wherein the microspheres of each subpopulation each comprise at least first and second target analytes attached to said microspheres with first and second attachment moieties, respectively;

wherein said microspheres are randomly distributed on said surface;

- 10
- b) contacting said array composition with a first set of extension probes that hybridize with at least said first target sequence adjacent to a first detection position to form an extension complex;

- c) contacting said extension complex with a composition comprising

- i) at least a first nucleotide;

- ii) polymerase;

wherein said polymerase extends a first extension probe with said first nucleotide when said first nucleotide is complementary to said first detection position of said first target sequence; and

- c) detecting the presence of said first nucleotide.

15

22 The method according to claim 21, wherein said first nucleotide comprises a label.

20

23. A method of determining the identification of a nucleotide at a detection position in at least a first target sequence comprising:

- 25
- a) providing an array composition comprising:

- i) a substrate with a surface comprising discrete sites; and

- ii) a population of microspheres comprising at least a first and a second subpopulation, wherein the microspheres of each subpopulation each comprise a plurality of target sequences;

- b) forming a first hybridization complex between said first target sequence and at least a first readout probe; and

- c) determining the nucleotide at said detection position.

30

24. A method according to claim 23, wherein said target sequence comprises a first and a second target domain, wherein said first hybridization complex comprises said first target sequence, a first readout probe hybridized to said first domain and a second readout probe hybridized to said second domain, wherein at least one of said readout probes comprise a label said determining comprises adding a ligase to form a ligation complex.

According to the nucleotide sequence

Add B2

5

gubB1

**SECRET**